

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number  
**WO 00/76540 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K 39/00**
- (21) International Application Number: **PCT/US00/15925**
- (22) International Filing Date: **9 June 2000 (09.06.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
**60/138,453** **10 June 1999 (10.06.1999)** **US**
- (71) Applicant: **MED IMMUNE, INC. [US/US];** 35 West Watkins Mill Road, Gaithersburg, MD 20878 (US).
- (72) Inventors: **ADAMOU, John, E.;** 20822 Shamrock Glen Circle, Germantown, MD 20874 (US). **CHOI, Gil, H.;** 11429 Potomac Oaks Drive, Rockville, MD 20850 (US).
- (81) Designated States (*national*): **AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (*regional*): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:  
— *Without international search report and to be republished upon receipt of that report.*
- (74) Agents: **GRANT, Alan, J. et al.;** Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **STREPTOCOCCUS PNEUMONIAE PROTEINS AND VACCINES**



(57) Abstract: The present invention relates to novel immunogenic polypeptides, and fragments thereof, and vaccines for the prevention and treatment of pneumococcal infection, especially by *Streptococcus pneumoniae*. The invention also relates to antibodies against the disclosed polypeptides, as well as vaccines containing said polypeptides and methods of disease prevention.

WO 00/76540 A2

## ***STREPTOCOCCUS PNEUMONIAE* PROTEINS AND VACCINES**

5

This application claims priority of U.S. provisional application 60/138,453, filed June 10, 1999, the disclosure of which is hereby incorporated by reference in its entirety.

10

### **BACKGROUND OF THE INVENTION**

15

*Streptococcus pneumoniae* is a gram positive bacterium which is a major causative agent in invasive infections in animals and humans, such as sepsis, meningitis, otitis media and lobar pneumonia (Tuomanen, et al. NEJM 322:1280-1284 (1995)). As part of the infective process, pneumococci readily bind to non-inflamed human epithelial cells of the upper and lower respiratory tract by binding to eukaryotic carbohydrates in a lectin-like manner (Cundell et al., Micro. Path. 17:361-374 (1994)). Conversion to invasive pneumococcal infections for bound bacteria may involve the local generation of inflammatory factors which may activate the epithelial cells to change the number and type of receptors on their surface (Cundell, et al., Nature, 377:435-438 (1995)). Apparently, one such receptor, platelet activating factor (PAF) is engaged by the pneumococcal bacteria and within a very short period of time (minutes) from the appearance of PAF, pneumococci exhibit strongly enhanced adherence and invasion of tissue. Certain soluble receptor analogs have

20

25

30

been shown to prevent the progression of pneumococcal infections (Idanpaan-Heikkila et al., J. Inf. Dis., 176:704-712 (1997)). A number of various other proteins have been suggested as being involved in the pathogenicity of *S. pneumoniae* but only some have been confirmed as virulence factors. Despite the fact that there are capsule conjugates currently in trial, there still remains a need for identifying additional polypeptides having epitopes in common from various strains of *S. pneumoniae* in order to utilize such polypeptides as vaccines to provide protection against a wide variety of *S. pneumoniae* serotypes.

15

#### BRIEF SUMMARY OF THE INVENTION

The invention disclosed herein relates to vaccines derived from polypeptides of the pneumococcal organism *Streptococcus pneumoniae*. In accordance with the present invention there are disclosed herein several protein sequences, and fragments thereof and their corresponding nucleotide sequences used for recombinantly preparing said polypeptides.

More specifically, the present invention discloses 2 large polypeptides, one denoted Sp128 (SEQ ID NO:6), composed of 664 amino acid residues, and a second polypeptide, denoted Sp130, containing 773 amino acid residues (SEQ ID NO:8). Both Sp128 and Sp130 have been found to confer protective properties on animals immunized with said polypeptides, or portions thereof.

The present invention also relates to the field of bacterial antigens and their use, for example, as immunogenic agents in humans and animals to stimulate an immune response. More specifically, it relates to the vaccination of mammalian species with one or more recombinant polypeptides produced according to the invention disclosed herein, such recombinant polypeptides being derived from *Streptococcus pneumoniae*.

In accordance with the present invention, such proteins serve as a mechanism for stimulating production of antibodies that protect the vaccine recipient against infection by a wide range of capsular serotypes of pathogenic *S. pneumoniae*.

The invention disclosed herein further relates to antisera and antibodies against such polypeptides useful in diagnosis and passive immune therapy with respect to diagnosing and treating such pneumococcal infections. Like the vaccines disclosed herein, the antibodies specific for such antigenic polypeptides, and fragments thereof, can be prepared recombinantly by transforming cells with vectors containing the appropriate gene sequences to produce the active tetrameric antibody. Such methods are well known in the art.

In a particular aspect, the present invention relates to the prevention and treatment of pneumococcal infections such as infections of the middle ear, nasopharynx, lung and bronchial areas, blood, CSF, and the like, that are caused by pneumococcal bacteria.

The present invention further relates to vaccines prepared from the novel proteins and polypeptides, as well

as fragments and segments thereof, disclosed herein. In addition, examples of the use of such proteins and polypeptides as vaccines for the protection of mammals are likewise disclosed.

5

#### BRIEF DESCRIPTION OF THE DRAWINGS

10

Figure 1 shows the results of 2 experiments (Figs 1A and 1B, respectively), using the same preparations of Sp128 and Sp130 polypeptides. The results demonstrate that active immunization with recombinant Sp128 or Sp130 polypeptides derived from the pneumococcal strain Norway serotype 4 is able to protect mice from death in a model of pneumococcal sepsis using the heterologous strain SJ2 (serotype 6B). In these 2 experiments, 90% and 100%, respectively, of the mice immunized with Sp130 survived the 14 day observation period following challenge with about 400 CFU (colony forming units) of pneumococci. Conversely, 100% of sham immunized mice (injected only with PBS (phosphate-buffered saline) plus adjuvant) died during the same period. In addition, for both experiments, 90% of the mice immunized with Sp128 survived the same 14 day observation period.

Figure 2 shows the results of passive administration of rabbit antiserum raised against Sp130 derived from Norway serotype 4. Such administration was able to protect mice in the pneumococcal sepsis model using a heterologous strain. More specifically, 70% of the mice immunized with the Sp130 antiserum survived the 10 day observation period after challenge with 1400 CFU of strain WU2 (serotype 3).

In addition, 100% of the mice immunized with a control serum (collected before immunization) died by day 4.

Figure 3 is a western blot showing reactivity of antisera raised against recombinant Sp130 (derived from strain Norway serotype 4) with whole cell lysates of heterologous strains. All *S. pneumoniae* strains tested showed a band of molecular weight about 220 kD, the expected mass for a protein containing both the Sp128 and Sp130 sequences, indicating that this protein was present in all of the tested strains. Tested strains included isolates from each of the pneumococcal serotypes represented in the currently used 23-valent polysaccharide vaccine.

15

Figure 4 is a western blot showing the reactivity of patient sera with either Sp128 or Sp130. Fig. 4A shows the results for Sp128. Fig. 4B shows the results for Sp130. The recombinant proteins were resolved by SDS-PAGE and transferred to nitrocellulose. Sera were collected from 5 patients (indicated by number at the top) at two different times. First collection (denoted "A" for "acute serum") was soon after onset of illness; second collection (denoted "C" for "convalescent") was made 8 to 30 days later. These sera were used to probe the blots. The results show that for patients 2, 3 and 5, convalescent serum reacted more strongly with Sp128 and Sp130 than did the corresponding acute serum. Such findings constitute indirect evidence that both Sp128 and Sp130 are expressed by *S. pneumoniae* during this phase of infection.

30

## DETAILED SUMMARY OF THE INVENTION

In accordance with the present invention there is disclosed herein recombinant polypeptides corresponding to  
5 Sp128 (SEQ ID NO: 6) and Sp130 (SEQ ID NO: 8).

It is an object of the present invention to provide methods of utilizing these recombinant polypeptides, and immunogenically active fragments thereof, as a means of  
10 immunizing animals, especially mammals, most especially humans, against a variety of microbial infections, especially pneumococcal infections.

It is a further object of the present invention to  
15 provide polypeptides, as disclosed herein, and active fragments thereof, whether derived from natural sources or prepared by means of recombinant technology, for use in immunizing animals, especially mammals, most especially humans, against pneumococcal infection.

20

It is a still further object of the present invention to provide vaccines that include polypeptides obtained from *S. pneumoniae* and/or variants of said polypeptides and/or active fragments of such polypeptides, including  
25 polypeptides prepared by recombinant means (i.e., recombinant polypeptides and proteins).

In accordance with the present invention, there are also disclosed herein nucleic acids and DNA sequences and  
30 molecules, and fragments thereof (and their corresponding isolated RNA sequences, and molecules and fragments thereof) showing sequence homology with, or identity to, or capable of hybridizing to, the DNA sequences identified in SEQ ID NOS: 5 and 7. The present invention

also relates to DNA (or RNA) sequences encoding the same polypeptide as is encoded by the sequences of SEQ ID NOS: 5 and 7, including fragments and portions thereof and, when derived from natural sources, including alleles thereof, for the express purpose of facilitating the recombinant expression of the immunogenic polypeptides, and immunogenic fragments thereof, disclosed herein.

Thus, an isolated DNA (or RNA) sequence can include only the coding region of the expressed gene (or fragment or portion thereof as hereinabove indicated) or can further include all or a portion of the non-coding DNA (or RNA) of the expressed human gene.

In accordance with the present invention, the term "percent identity" or "percent identical," when referring to a sequence, means that a sequence is compared to a claimed or described sequence after alignment of the sequence to be compared (the "Compared Sequence") with the described or claimed sequence (the "Reference Sequence"). The Percent Identity is then determined according to the following formula:

$$\text{Percent Identity} = 100 [1 - (C/R)]$$

25

wherein C is the number of differences between the Reference Sequence and the Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence wherein (i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence and (ii) each gap in the Reference Sequence and (iii) each aligned base or amino acid in the Reference Sequence that



is different from an aligned base or amino acid in the Compared Sequence, constitutes a difference; and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

If an alignment exists between the Compared Sequence and the Reference Sequence for which the percent identity as calculated above is about equal to or greater than a specified minimum Percent Identity then the Compared Sequence has the specified minimum percent identity to the Reference Sequence even though alignments may exist in which the hereinabove calculated Percent Identity is less than the specified Percent Identity.

In accordance with the present invention, there are disclosed herein the polynucleotide sequences coding for the polypeptide vaccines of the invention so as to facilitate recombinant expression of said polypeptides. Such polynucleotides code for the polypeptides of SEQ ID NOS: 6 and 8 and are disclosed as the sequences of SEQ ID NOS: 5 and 7.

For purposes of recombinantly expressing the polypeptide vaccines of the invention, the polynucleotides of SEQ ID NOS: 5 and 7 may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexa-histidine tag (for example, as can be supplied by a pQE-9 vector) to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a

mammalian host, e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

5

To facilitate generation of the polynucleotides disclosed herein, appropriate PCR primers are provided as SEQ ID NOS: 1 (5'-primer for Sp128), 2 (3'-primer for Sp128), 3 (5'-primer for Sp130), and 4 (3'-primer for Sp130).

10

The polypeptides, and fragments thereof, of the vaccines disclosed as expression products according to the invention may be in "enriched form." As used herein, the term "enriched" means that the concentration of the material is at least about 2, 5, 10, 100, or 1000 times its natural concentration (for example), advantageously 0.01%, by weight, preferably at least about 0.1% by weight. Enriched preparations of about 0.5%, 1%, 5%, 10%, and 20% by weight are also contemplated. The sequences, constructs, vectors, clones, and other materials comprising the present invention can advantageously be in enriched or isolated form.

25

"Isolated" in the context of the present invention with respect to polypeptides means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living organism is not isolated, but the same polypeptide, separated from some or all of the co-existing materials in the natural system, is isolated. Such polypeptides could be part of a composition, and still be isolated in that such composition is not part of its

30

natural environment. The polypeptides of the vaccines disclosed herein are preferably provided in an isolated form, and preferably are purified to homogeneity.

5       The recombinant or immunogenic polypeptides disclosed in accordance with the present invention may also be in "purified" form. The term "purified" does not require absolute purity; rather, it is intended as a relative definition, and can include preparations that  
10   are highly purified or preparations that are only partially purified, as those terms are understood by those of skill in the relevant art. For example, individual clones isolated from a cDNA library have been conventionally purified to electrophoretic homogeneity.  
15   Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. Furthermore, claimed polypeptide which has a purity of preferably  
20   0.001%, or at least 0.01% or 0.1%; and even desirably 1% by weight or greater is expressly contemplated.

      The term "coding region" refers to that portion of a gene which either naturally or normally codes for the  
25   expression product of that gene in its natural genomic environment, i.e., the region coding *in vivo* for the native expression product of the gene. The coding region can be from a normal, mutated or altered gene, or can even be from a DNA sequence, or gene, wholly synthesized  
30   in the laboratory using methods well known to those of skill in the art of DNA synthesis.

      The term "primer" means a short nucleic acid sequence that is paired with one strand of DNA and

provides a free 3'OH end at which a DNA polymerase starts synthesis of a deoxyribonucleotide chain.

At the simplest level, the amino acid sequence  
5 corresponding to all or part of the polypeptides  
according to the present invention can be synthesized  
using commercially available peptide synthesizers. This  
is particularly useful in producing small peptides and  
fragments of larger polypeptides. (Fragments are useful,  
10 for example, in generating antibodies against the native  
polypeptide.)

The terms "fragment," "derivative" and "analog," when  
referring to the polypeptides according to the present  
15 invention, means a polypeptide which retains essentially  
the same biological function or activity as said  
polypeptide. Thus, an analog includes a proprotein which  
can be activated by cleavage of the proprotein portion to  
produce an active mature polypeptide. Such fragments,  
20 derivatives and analogs must have sufficient similarity to  
the polypeptides SEQ ID NOS: 6 and 8, so that activity of  
the native polypeptide is retained.

The polypeptide vaccines of the present invention may  
25 be recombinant polypeptides, natural polypeptides or  
synthetic polypeptides, preferably recombinant  
polypeptides.

"Recombinant," as used herein, means that a protein  
30 is derived from recombinant (e.g., microbial or  
mammalian) expression systems. "Microbial" refers to  
recombinant proteins made in bacterial or fungal (e.g.,  
yeast) expression systems. As a product, "recombinant  
microbial" defines a protein essentially free of native

endogenous substances and unaccompanied by associated native glycosylation. Protein expressed in most bacterial cultures, e.g., E. coli, will be free of glycosylation modifications that might normally occur in yeast or mammalian expression systems. Thus, the patterns of such post-translational modifications will differ with the expression system. However, all such variants are considered to lie within the disclosure of the present invention.

A vaccine according to the present invention would include a polypeptide, including immunogenic fragments thereof, comprising an amino acid sequence at least 65% identical, preferably 80% identical, most preferably 95% identical and ideally 100% identical to the amino acid sequence of SEQ ID NO:6.

Such vaccines would also comprise a polypeptide, including immunogenic fragments thereof, having an amino acid sequence at least 65% identical, preferably 80% identical, most preferably 95% identical, and ideally 100% identical to the amino acid sequence of SEQ ID NO:8.

The present invention is also directed to an antiserum produced by immunizing an animal with a polypeptide according to the invention. The invention also includes and isolated antibody that binds specifically to a polypeptide of the invention. Such an antibody may be a monoclonal antibody, possibly produced by a hybridoma cell line, and may also include a recombinantly produced antibody formed by introducing into a suitable cell line the gene sequences required for producing an antibody specific for the polypeptide vaccines disclosed herein.

The present invention is also directed to a vaccine comprising one or more *S. pneumoniae* polypeptides selected from the polypeptides, and immunogenic fragments thereof, disclosed herein, suspended in a pharmaceutically acceptable diluent, carrier or excipient, provided that said polypeptide is present in an amount effective to elicit protective antibodies in an animal against an organism related to the genus *Streptococcus*, preferably an organism of the genus *Streptococcus*, and most preferably where the organism is *Streptococcus pneumoniae*.

The present invention also provides for a method of preventing or treating an infection caused by a member of the genus *Streptococcus* in an animal, comprising administering to an animal, especially a mammal, and most especially a human being, a polypeptide, or immunogenic fragment thereof, as disclosed herein, and wherein said polypeptide, or immunogenic fragment thereof, is administered in an amount effective to prevent or attenuate said infection. In using the methods of the invention, the disease to be prevented or treated will preferably be a pneumococcal infection, most preferably an infection by an organism that is a member of the genus *Streptococcus*, ideally *Streptococcus pneumoniae*.

A vaccine disclosed according to the present invention may also include a vaccine comprising a microbial organism transformed with polynucleotides, and thereby expressing the polypeptides, or fragments thereof, selected from the group consisting of Sp128 and Sp130 (SEQ ID NOS: 6 and 8, respectively). The present invention would thus also encompass a method of preventing or attenuating an infection caused by a member

of the genus *Streptococcus* in an animal, especially a mammal, most especially a human, comprising administering to said animal such a vaccine, wherein said vaccine is administered in an amount effective to prevent or  
5 attenuate said infection. In applying the method of the invention, the transformed microorganism is selected from the group consisting of *Salmonella*, *Mycobacteria*, *Streptococcus*, poxviruses, and adenoviruses.

10 Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides.  
15 Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

The immunogenic fragments of the polypeptide vaccines  
20 disclosed according to the invention will include immunogenic fragments of Sp128 (SEQ ID NO:6), which fragments can be readily screened for immunogenic activity, as well as immunogenic fragments of Sp130 (SEQ ID NO: 8). For example, in the amino acid sequence of  
25 Sp130, the fragment corresponding to residues 657 through 773 are known to provide about 40% protection versus the entire Sp130 sequence. Thus, the former fragment protects about 4 out of 10 mice challenged with *Streptococcus pneumoniae* versus 10 of 10 for the entire Sp130 sequence.  
30 Thus, specific fragment may include the fragments having amino acid sequences 650 - 773, 640 - 773, 630 - 773, 620 - 773, 610 - 773, 600 - 773, and similar fragments up to the entire Sp130 sequence (SEQ ID NO: 8). It is logical to presume that fragments of Sp128 (SEQ ID NO: 6) may provide

similar degree of protection versus the entire Sp128 protein.

Such variations in homology for putative vaccines are well known in the art (See, for example, Hansen et al., "Active and Passive Immunity Against *Borelia bergdorferi* Decorin Binding Protein A (DbpA)," Infection and Immunity, (May) 1998, p. 2143 - 2153; Roberts et al., "Heterogeneity Among Genes Including Decorin Binding Proteins A and B of *Borelia bergdorferi sensu lato*," Infection and Immunity, (Nov) 1998, p. 5275-5285). Such observations would similarly apply to portions of the proteins disclosed herein.

Such fragments or segments find a multitude of uses. For example, such segments of the polypeptides according to the present invention find use as intermediates in the synthesis of higher molecular weight structures also within the present invention.

The term "active fragment" or "immunogenic fragment" means a fragment that generates an immune response (i.e., has immunogenic activity) when administered, alone or optionally with a suitable adjuvant, to an animal, such as a mammal, for example, a rabbit or a mouse, and also including a human.

As noted, the polypeptides, fragments or other derivatives, or analogs thereof, or cells expressing them, can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal, monoclonal, chimeric, single chain, Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the



production of polyclonal antibodies, especially where these are in the form of antisera raised against the polypeptides, or fragments thereof, according to the present invention. Such antisera find use in immunization  
5 against pneumococcal infection.

Antibodies generated against a polypeptide vaccine corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptide into  
10 an animal or by administering the polypeptide to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptide itself. In this manner, even a sequence encoding only a fragment of the polypeptide can be used to generate antibodies binding  
15 the whole native polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples  
20 include the hybridoma technique (Kohler and Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al.,  
25 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

Thus, the present invention also relates to the use of the novel polypeptides disclosed herein, as well as to  
30 immunogenic fragments thereof, for the production of lymphocytes, or hybridoma cells, producing monoclonal antibodies against such polypeptides, or immunogenic fragments thereof. The present invention also relates to the hybridoma cells producing such antibodies.

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention.

The antibodies can be used in methods relating to the localization and activity of the protein sequences of the invention, e.g., for imaging these proteins, measuring levels thereof in appropriate physiological samples and the like, and for other diagnostic applications.

A vaccine in accordance with the present invention may include one or more of the hereinabove described polypeptides or active fragments thereof. When employing more than one polypeptide or active fragment, such as two or more polypeptides and/or active fragments may be used as a physical mixture or as a fusion of two or more polypeptides or active fragments. The fusion fragment or fusion polypeptide may be produced, for example, by recombinant techniques or by the use of appropriate linkers for fusing previously prepared polypeptides or active fragments.

25

In many cases, a variation in the polypeptide or active fragment is a conservative amino acid substitution, although other substitutions are within the scope of the invention.

30

In accordance with the present invention, a polypeptide variant includes variants in which one or more amino acids are substituted and/or deleted and/or inserted.

In another aspect, the invention relates to passive immunity vaccines formulated from antibodies against a polypeptide or active fragment of a polypeptide of the present invention. Such passive immunity vaccines can be utilized to prevent and/or treat pneumococcal infections in patients. In this manner, according to a further aspect of the invention, a vaccine can be produced from a synthetic or recombinant polypeptide of the present invention or an antibody against such polypeptide.

As already described, another aspect the present invention relates to a method of using one or more antibodies (monoclonal, polyclonal or sera) to the polypeptides of the invention as described above for the prophylaxis and/or treatment of diseases that are caused by pneumococcal bacteria. In particular, the invention relates to a method for the prophylaxis and/or treatment of infectious diseases that are caused by *S. pneumoniae*. In a still further preferred aspect, the invention relates to a method for the prophylaxis and/or treatment of otitis media, nasopharyngeal and bronchial infections, and the like in humans by utilizing a vaccine of the present invention.

25

Generally, vaccines are prepared as injectables, in the form of aqueous solutions or suspensions. Vaccines in an oil base are also well known such as for inhaling. Solid forms which are dissolved or suspended prior to use may also be formulated. Pharmaceutical carriers, diluents and excipients are generally added that are compatible with the active ingredients and acceptable for pharmaceutical use. Examples of such carriers include, but are not limited to, water, saline solutions, dextrose,

30

or glycerol. Combinations of carriers may also be used.

Vaccine compositions may further incorporate additional substances to stabilize pH, or to function as  
5 adjuvants, wetting agents, or emulsifying agents, which can serve to improve the effectiveness of the vaccine.

Vaccines are generally formulated for parenteral administration and are injected either subcutaneously or  
10 intramuscularly. Such vaccines can also be formulated as suppositories or for oral administration, using methods known in the art, or for administration through nasal or respiratory routes.

15 The amount of vaccine sufficient to confer immunity to pathogenic bacteria is determined by methods well known to those skilled in the art. This quantity will be determined based upon the characteristics of the vaccine recipient and the level of immunity required. Typically,  
20 the amount of vaccine to be administered will be determined based upon the judgment of a skilled physician. Where vaccines are administered by subcutaneous or intramuscular injection, a range of .5 to 500  $\mu$ g purified protein may be given.

25 The present invention is also directed to a vaccine in which a polypeptide or active fragment of the present invention is delivered or administered in the form of a polynucleotide encoding the polypeptide or active  
30 fragment, whereby the polypeptide or active fragment is produced in vivo. The polynucleotide may be included in a suitable expression vector and combined with a pharmaceutically acceptable carrier.

In addition, the polypeptides of the present invention can be used as immunogens to stimulate the production of antibodies for use in passive immunotherapy, for use as diagnostic reagents, and for use as reagents in other processes such as affinity chromatography.

In another aspect the present invention provides polynucleotides which encode the hereinabove described polypeptides and active fragments of the invention. The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors comprising a polynucleotide encoding a polypeptide of the invention. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the polynucleotides which encode such polypeptides. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

Vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other

vector may be used as long as it is replicable and viable in the host.

5 The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

10 The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the E. coli lac or trp,  
15 the phage lambda P<sub>L</sub> promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation  
20 initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably  
25 contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

30 The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the

proteins.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli,  
5 Streptomyces, Salmonella typhimurium; fungal cells, such as yeast; insect cells such as Drosophila S2 and Spodoptera Sf9; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of  
10 those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs  
15 comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter,  
20 operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example. Bacterial: pQE70, pQE60, pQE-9 (Qiagen, Inc.), pBS, pD10, phagescript, psiX174,  
25 pbluescript SK, pBS, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long  
30 as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate

vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P<sub>R</sub>, P<sub>L</sub> and TRP. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), the



disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of E. coli and S. cerevisiae TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK),  $\alpha$ -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

30

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable

reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice, including streptococcal species, especially *S. pneumoniae*.

In a further embodiment, microbial organisms genetically transformed with polynucleotides expressing Sp128 or Sp130, or both, may themselves be used as living vaccine delivery vehicles. Examples include, but are in no way limited to, *Salmonella* species, *Mycobacterium* species, *Streptococcus* species, poxviruses, adenoviruses, and the like. In addition, transgenic edible plants may also be candidates for vaccine delivery.

20

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Amersham Pharmacia Biotech, Piscataway, NJ, USA) and pGEM1 (Promega, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

30

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell

density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

- 5        Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

10        Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, a french press, mechanical disruption, or use of cell lysing agents, such methods are well know to those skilled in the art. However, preferred are host cells which secrete the  
15        polypeptide of the invention and permit recovery of the polypeptide from the culture media.

20        Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will  
25        comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the  
30        SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The polypeptides can be recovered and/or purified from recombinant cell cultures by well-known protein

recovery and purification methods. Such methodology may include ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. In this respect, chaperones may be used in such a refolding procedure. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides that are useful as immunogens in the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated.

Procedures for the isolation of the individually expressed polypeptides may be isolated by recombinant expression/isolation methods that are well-known in the art. Typical examples for such isolation may utilize an antibody to a conserved area of the protein or to a His tag or cleavable leader or tail that is expressed as part of the protein structure.

30

Specific embodiments of the invention will now be further described in more detail in the following non-limiting examples and it will be appreciated that additional and different embodiments of the teachings of

the present invention will doubtless suggest themselves to those of skill in the art and such other embodiments are considered to have been inferred from the disclosure herein.

5

### EXAMPLE 1

#### Active Protection with Anti-Sp128 and Anti-Sp130

10

A. Cloning, Expression, and Purification of Sp128 and Sp130.

The genomic DNA used as target for amplification using the polymerase chain reaction was isolated from  
15 *Streptococcus pneumoniae* (Norway strain - serotype 4), the same strain used for genomic sequencing. The nucleotide sequence of the gene fragments encoding Sp128 (SEQ ID NO: 5) and Sp130 (SEQ ID NO: 7) with the corresponding amino acid sequence for polypeptides Sp128  
20 (SEQ ID NO: 6) and Sp130 (SEQ ID NO: 8) are provided in the Sequence Listing.

Primers (SEQ ID NOS: 1 - 4) were designed so as to amplify either Sp128 or Sp130 gene fragments and allow  
25 their cloning into the *E. coli* expresssion vector pQE10 with, for example, subsequent expression of a histidine-tagged protein product for purification by Nickel-affinity chromatography.

30 Thus, cloning of the fragments amplified by the primers of SEQ ID NOS: 1 and 2 results in the polypeptide of SEQ ID NO: 6 (denoted Sp128), while cloning of the fragment amplified using the primers disclosed in SEQ ID NOS: 3 and 4 result in the polypeptide of SEQ ID NO: 8

(denoted Sp130).

5 B. Vaccination With Sp128 and Sp130 Results in Protection  
Against Lethal *S. pneumoniae* Challenge

10 In each of the experiments shown in Figure 1,  
C3H/HeJ mice (10 per group) were immunized  
intraperitoneally (i.p.) with either Sp128 or Sp130  
protein (15 µg in 50 µl PBS (phosphate buffered saline)  
emulsified in 50 µl complete Freund's adjuvant (CFA)). A  
group of 10 mice were sham-immunized with PBS and CFA  
only.

15

A second immunization of 15 µg protein with  
incomplete Freund's adjuvant (IFA) was administered 3  
weeks later (with the sham-immunized group receiving PBS  
and IFA).

20

Blood was drawn (retro-orbital bleed) at week 7.  
Sera from each group were pooled for analysis of anti-  
Sp128 and anti-Sp130 antibody by ELISA. Mice were  
challenged at week 8 by intraperitoneal injection of  
25 approximately 400 CFU (colony forming units) of *S.*  
*pneumoniae* strain SJ2 (capsular serotype 6B). In  
preliminary experiments, the median infection dose  
(LD<sub>50</sub>) of this strain was determined to be approximately 10  
CFU. Mice were monitored for 14 days of survival.

30

Both experiments shown in Figure 1 used the same  
preparations of recombinant Sp128 and Sp130.

In the experiment shown in Figure 1A, 7-week serum

collected from the 10 mice immunized with either Sp128 or Sp130 each had an endpoint ELISA titer of 1:2,048,000 and 1:1,024,000, respectively. No anti-Sp128 or anti-Sp130 antibody was detected in sera from sham-immunized mice.

5 Ninety percent of the mice immunized with either Sp128 or Sp130 protein survived the challenge (406 CFU of pneumococci) for the extent of the study (14 days). One hundred percent of sham-immunized mice were dead by day 7.

10

In the experiment shown in Figure 1B, 7-week sera collected from the 10 mice immunized with either Sp128 or Sp130 each had an endpoint ELISA titer of 1:1,024,000 and 1:512,000, respectively. No anti-Sp128 or anti-Sp130 antibody was detected in sera from sham-immunized mice. Ninety and one hundred percent of the mice immunized with either Sp128 or Sp130 protein, respectively, survived the challenge (404 CFU of pneumococci) for the extent of the study (14 days). One hundred percent of sham-immunized mice were dead by day 5.

20

These data indicate that immunization of mice with either recombinant Sp128 or Sp130 proteins elicit a response capable of protecting against systemic pneumococcal infection and subsequent death. Cross protection is demonstrated by the fact that the recombinant pneumococcal protein was generated based on capsular serotype 4 DNA sequence, while the challenge was with the heterologous strain SJ2 (capsular serotype 6B).

30

## EXAMPLE 2

### Passive Protection With Anti-Sp130 Antisera

#### A. Generation of Rabbit Immune Sera

5           Following collection of pre-immune serum, a New Zealand White rabbit was immunized with 250 µg of Sp130 (SEQ ID NO:8) in complete Freund's adjuvant. The rabbit was given 2 boosts of 125 µg Sp130 in incomplete Freund's  
10   adjuvant on days 21 and 52, and bled on days 31 and 62.

#### B. Passive Protection in Mice

15           BALB/cByJ mice (10 per group) were passively immunized by 2 i.p. injections of 100 µl of rabbit serum. The first injection was administered 24 hours before challenge with 1400 CFU of *S. pneumoniae* strain WU2, and the second injection was given 4 hours after challenge.  
20   Figure 2 shows the survival of mice after infection with WU2 (capsular serotype 3) strain. In preliminary experiments, the LD<sub>50</sub> of this strain was determined to be approximately 100 CFU.

25           Figure 2 shows the survival of mice injected with 1400 CFU of strain WU2. As shown therein, 70% of the mice immunized with rabbit immune serum raised against Sp130 protein survived the 10 day observation period. Of the mice immunized with the control serum (collected from a  
30   rabbit prior to immunization), 100% died by day 4.

          These data suggest that the protection against pneumococcal infection resulting from immunization with Sp130 is antibody-mediated, since the mice were protected



by passive transfer of serum from a hyperimmunized rabbit. As seen in the previously described mouse active challenge experiments, serum directed against recombinant Sp130 protein cloned from a serotype 4 strain was protective against challenge with a heterologous strain, WU2 (capsular serotype 3).

10

15

### EXAMPLE 3

Conservation of Sp128-Sp130 Among Strains of S.  
pneumoniae

#### A. Western Blotting

The pneumococcal strains used in this experiment were obtained from the American Type Culture Collection (Rockville, MD) and include one isolate from each of the serotypes in the currently used multivalent pneumococcal vaccine.

For total cell lysates, pneumocci were grown to mid-logarithmic phase (absorbance at 620 nm was 0.4 to 0.6) in 2 ml Todd-Hewitt broth with 5% yeast extract (from Difco, Detroit, MI) at 37°C. Bacteria were harvested by centrifugation and washed twice with water. Pellets

(consisting of sedimented cells) were resuspended in 200  $\mu$ l of lysis buffer (0.01% sodium dodecyl sulfate, 0.15 M sodium citrate, and 0.1% sodium deoxycholate) and incubated at 37°C for 30 min, then diluted in an equal volume of 2X SSC (0.3 M NaCl, 0.03 M sodium citrate). Polypeptides in the lysates were resolved on SDS-PAGE (sodium dodecyl sulfate polyacrylamide gel electrophoresis), transferred to nitrocellulose membranes (Bio-Rad Laboratories, Hercules, CA) and probed with antibody by conventional Western Blotting procedures. Sera from a New Zealand White rabbit immunized with Sp130 (as per Example 2, supra) was used at a dilution of 1:3000. Bound antibody was detected with peroxidase-conjugated sheep anti-rabbit IgG using a chemiluminescence kit from Amersham Inc. (Cambridge, MA).

The rabbit anti-Sp130 sera revealed 2 major bands with apparent molecular weights of 110 kD and 220 kD in all 23 pneumococcal lysates tested (as shown in Figure 3).

These data show that Sp130 is expressed and shares common antigenic epitopes among strains of the 23 pneumococcal capsular serotypes represented in the currently used polysaccharide vaccine.

30

#### EXAMPLE 4

##### Immunogenicity of Sp128 and Sp130 in Humans

Sera from patients with culture-proven pneumococcal

bacteremia were used in Western blots containing recombinant Sp128 or Sp130 protein. In the experiment shown in Figure 4, sera from 5 patients (indicated by numerals 1 through 5) were diluted 1:3000 and used to probe blots containing Sp128 (SEQ ID NO:6) or Sp130 (SEQ ID NO:8).

The lanes labeled "A" (for "acute") were probed with serum collected shortly after diagnosis of pneumococcal infection; lanes denoted "C" (for "convalescent") were probed with serum collected either 1 month later (patients 1, 2, and 3) or 8 days after the first serum collection (patients 4 and 5). For patients 2, 3, and 5, reactivity of the convalescent serum with Sp128 and Sp130 was stronger than that of the corresponding acute serum.

Other experiments (not depicted in the figure) showed that convalescent sera from 17 patients with pneumococcal infections were tested individually for reactivity with either Sp128 or Sp130. Thus, 10 and 15 of the 23 sera were found to bind (on a Western Blot) Sp128 and Sp130, respectively.

These experiments indicate that Sp128 and Sp130 (the latter to a greater extent), are recognized by the human immune system and suggest that antibodies able to bind the Sp128-Sp130 protein may be produced during natural *S. pneumoniae* infection in humans. Further, this provides indirect evidence that Sp128 and Sp130 are expressed in vivo by *S. pneumoniae* during this phase of infection, and thus may be available as targets for immunoprophylaxis, immunotherapy, or to provide anamnestic immune responses in subjects vaccinated with these proteins. Since the patients were infected with a variety of pneumococcal

strains, these data also support the idea that Sp130 is more antigenically conserved than Sp128.

5

10

15

20

25

30

WHAT IS CLAIMED IS:

1. A vaccine comprising a polypeptide, including immunogenic fragments thereof, having an amino acid sequence at least 65% identical to the amino acid sequence of SEQ ID NO:6.  
5
2. The vaccine of claim 1 wherein said amino acid sequence is at least 80% identical to the amino acid sequence of SEQ ID NO:6.  
10
3. The vaccine of claim 1 wherein said amino acid sequence is at least 95% identical to the amino acid sequence of SEQ ID NO:6.  
15
4. The vaccine of claim 1 wherein said amino acid sequence is identical to the amino acid sequence of SEQ ID NO:6.
- 20 5. A vaccine comprising a polypeptide, including immunogenic fragments thereof, having an amino acid sequence at least 65% identical to the amino acid sequence of SEQ ID NO:8.
- 25 6. The vaccine of claim 5 wherein said amino acid sequence is at least 80% identical to the amino acid sequence of SEQ ID NO:8.
- 30 7. The vaccine of claim 5 wherein said amino acid sequence is at least 95% identical to the amino acid sequence of SEQ ID NO:8.
8. The vaccine of claim 5 wherein said amino acid sequence is identical to the amino acid sequence of SEQ

ID NO:8.

9. An antiserum produced by immunizing an animal with a polypeptide selected from the group consisting of the polypeptides of claims 1, 2, 3, 4, 5, 6, 7, and 8.

10. An isolated antibody that binds specifically to a polypeptide selected from the group consisting of the polypeptides according to claims 1, 2, 3, 4, 5, 6, 7, and 8.

11. The antibody of claim 10 wherein the antibody is a monoclonal antibody.

12. An engineered cell producing a monoclonal antibody of claim 11.

13. An antiserum produced by immunizing an animal with the polypeptide of SEQ ID NO: 6.

14. An antiserum produced by immunizing an animal with the polypeptide of SEQ ID NO: 8.

15. An isolated recombinant antibody that binds specifically to a polypeptide selected from the group consisting of the polypeptides of claims 1,2,3,4,5,6,7, and 8.

16. A vaccine comprising :  
a. one or more *S. pneumoniae* polypeptides selected from the group consisting of the polypeptides of claims 1, 2, 3, 4, 5, 6, 7, and 8; and  
b. a pharmaceutically acceptable diluent , carrier or excipient;

wherein said polypeptide is present in an amount effective to elicit protective antibodies in an animal against an organism of the genus *Streptococcus*.

5        17. A method of preventing or attenuating an infection caused by a member of the genus *Streptococcus* in an animal, comprising administering to said animal a polypeptide selected from the group consisting of the polypeptides of claims 1, 2, 3, 4, 5, 6, 7, and 8, and  
10        wherein said polypeptide is administered in an amount effective to prevent or attenuate said infection.

18. A method of preventing pneumococcal infection by administering to an animal the vaccine according to claim  
15        16.

19. A method of preventing or attenuating an infection caused by a member of the genus *Streptococcus* in an animal, comprising administering to said animal an  
20        antibody according to claim 10, wherein said antibody is administered in an amount effective to prevent or attenuate said infection.

20. A vaccine comprising a microbial organism  
25        transformed with polynucleotides, and thereby expressing the polypeptides, or fragments thereof, selected from the group consisting of Sp128 and Sp130.

21. A method of preventing or attenuating an  
30        infection caused by a member of the genus *Streptococcus* in an animal, comprising administering to said animal a vaccine according to claim 20, wherein said antibody is administered in an amount effective to prevent or attenuate said infection.

22. The vaccine according to claim 20, wherein said  
transformed microorganism is selected from the group  
consisting of Salmonella, Mycobacteria, Streptococcus,  
5 poxviruses, and adenoviruses.

10

15

20

25

30



FIG. 1A




FIG. 1B




FIG. 2






FIG. 3



## FIG. 4



## SEQUENCE LISTING

<110> Adamou, John  
Choi, Gil

<120> Streptococcus Pneumoniae Proteins and Vaccines

<130> 469201-475

<140>

<141>

<150> U.S. 60/138,453

<151> 1999-06-10

<160> 8

<170> PatentIn Ver. 2.1

<210> 1

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Forward primer  
for PCR amplification of Sp128 genomic sequences

<400> 1

tacccggtag tcttagcaga c

21

<210> 2

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Reverse primer  
for PCR amplification of Sp128 genomic sequence

<400> 2

atagccataa gttgatttgc catta

25

<210> 3

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Forward primer  
for PCR amplification of Sp130 genomic sequence

<400> 3

aagcttggcg agattgcaga a

21

<210> 4

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Reverse primer  
for PCR amplification of Sp130 genomic sequence

<400> 4

cttattagga ttgttagtag ttgattt

27

<210> 5

<211> 1992

<212> DNA

<213> Streptococcus pneumoniae

<400> 5

taccggtag tcttagcaga cacatctagc tctgaagatg ctttaaacaat ctctgataaa 60  
gaaaaagtag cagaaaataa agagaaacat gaaaatatcc atagtgcctat ggaaacttca 120  
caggatttta aagagaagaa aacagcagtc attaaggaaa aagaagttgt tagtaaaaaat 180  
cctgtgatag acaataaacac tagcaatgaa gaagcaaaaa tcaaagaaga aaattccaat 240  
aaatcccaag gagattatac ggactcattt gtgaataaaa acacagaaaa tccccaaaaa 300  
gaagataaag ttgtctatat tgctgaattt aaagataaag aatctggaga aaaagcaatc 360  
aaggaactat ccagtcttaa gaatacaaaa gttttatata cttatgatag aatttttaac 420  
ggtagtgcca tagaaacaac tccagataac ttggacaaaa ttaaacaat agaaggtatt 480  
tcatcggttg aaagggcaca aaaagtccaa cccatgatga atcatgccag aaaggaaatt 540  
ggagttgagg aagctattga ttacctaaag tctatcaatg ctccgtttg gaaaaatttt 600  
gatggtagag gtatggtcat ttcaaatac gatactggaa cagattatag acataaggct 660  
atgagaatcg atgatgatgc caaagcctca atgagattta aaaaagaaga cttaaaaggc 720  
actgataaaa attattgggt gagtgataaa atccctcatg cgttcaatta ttataatggg 780  
ggcaaatca ctgtagaaaa atatgatgat ggaagggatt attttgaccc acatgggatg 840  
catattgcag ggattcttgc tggaaatgat actgaacaag acatcaaaaa ctttaacggc 900  
atagatggaa ttgcacctaa tgcacaaatt ttctcttaca aaatgtattc tgacgcagga 960  
tctgggtttg cgggtgatga aacaatgttt catgctattg aagattctat caaacacaac 1020  
gttgatgttg tttcggtatc atctggtttt acaggaacag gtctttagg tgagaaatat 1080  
tggcaagcta ttcgggcatt aagaaaagca ggcattccaa tgggtgtcgc tacgggtaac 1140

tatgcgactt ctgcttcaag ttcttcatgg gatttagtag caaataatca tctgaaaaatg 1200  
 accgacactg gaaatgtaac acgaactgca gcacatgaag atgcatagc ggtcgcttct 1260  
 gctaaaaatc aaacagttga gtttgataaa gttaacatag gtggagaaag ttttaaatac 1320  
 agaaatatag gggccttttt cgataagagt aaaatcacaa caaatgaaga tggaacaaaa 1380  
 gctcctagta aattaaaatt tgtatatata ggcaaggggc aagaccaaga tttgataggt 1440  
 ttggatctta ggggcaaaat tgcagtaatg gatagaattt atacaaagga tttaaaaaat 1500  
 gcttttaaaa aagctatgga taagggtgca cgcgccatta tgggtgtaaa tactgtaaat 1560  
 tactacaata gagataattg gacagagctt ccagctatgg gatatgaagc ggatgaaggt 1620  
 actaaaagtc aagtgttttc aatttcagga gatgatgggtg taaagctatg gaacatgatt 1680  
 aatcctgata aaaaaactga agtcaaaaaga aataataaag aagattttta agataaattg 1740  
 gagcaatact atccaattga tatggaaagt ttttaattcca acaaaccgaa ttaggtgac 1800  
 gaaaaagaga ttgactttta gtttgcacct gacacagaca aagaactcta taaagaagat 1860  
 atcatcgctt cagcaggatc tacatcttgg gggccaagaa tagatttact tttaaaaccc 1920  
 gatgtttcag cacctggtaa aaatattaaa tccacgctta atgttattaa tggcaaatca 1980  
 acttatggct at 1992

<210> 6

<211> 664

<212> PRT

<213> Streptococcus pneumoniae

<400> 6

Tyr	Pro	Val	Val	Leu	Ala	Asp	Thr	Ser	Ser	Ser	Glu	Asp	Ala	Leu	Asn
1				5				10						15	
Ile	Ser	Asp	Lys	Glu	Lys	Val	Ala	Glu	Asn	Lys	Glu	Lys	His	Glu	Asn
			20					25					30		
Ile	His	Ser	Ala	Met	Glu	Thr	Ser	Gln	Asp	Phe	Lys	Glu	Lys	Lys	Thr
		35					40					45			
Ala	Val	Ile	Lys	Glu	Lys	Glu	Val	Val	Ser	Lys	Asn	Pro	Val	Ile	Asp
	50					55					60				
Asn	Asn	Thr	Ser	Asn	Glu	Glu	Ala	Lys	Ile	Lys	Glu	Glu	Asn	Ser	Asn
65					70					75				80	
Lys	Ser	Gln	Gly	Asp	Tyr	Thr	Asp	Ser	Phe	Val	Asn	Lys	Asn	Thr	Glu
			85						90					95	
Asn	Pro	Lys	Lys	Glu	Asp	Lys	Val	Val	Tyr	Ile	Ala	Glu	Phe	Lys	Asp
		100						105					110		
Lys	Glu	Ser	Gly	Glu	Lys	Ala	Ile	Lys	Glu	Leu	Ser	Ser	Leu	Lys	Asn
	115					120					125				
Thr	Lys	Val	Leu	Tyr	Thr	Tyr	Asp	Arg	Ile	Phe	Asn	Gly	Ser	Ala	Ile

130		135		140	
Glu Thr Thr Pro Asp Asn Leu Asp Lys Ile Lys Gln Ile Glu Gly Ile					
145		150		155	160
Ser Ser Val Glu Arg Ala Gln Lys Val Gln Pro Met Met Asn His Ala					
	165		170		175
Arg Lys Glu Ile Gly Val Glu Glu Ala Ile Asp Tyr Leu Lys Ser Ile					
	180		185		190
Asn Ala Pro Phe Gly Lys Asn Phe Asp Gly Arg Gly Met Val Ile Ser					
	195		200		205
Asn Ile Asp Thr Gly Thr Asp Tyr Arg His Lys Ala Met Arg Ile Asp					
	210		215		220
Asp Asp Ala Lys Ala Ser Met Arg Phe Lys Lys Glu Asp Leu Lys Gly					
	225		230		235
Thr Asp Lys Asn Tyr Trp Leu Ser Asp Lys Ile Pro His Ala Phe Asn					
	245		250		255
Tyr Tyr Asn Gly Gly Lys Ile Thr Val Glu Lys Tyr Asp Asp Gly Arg					
	260		265		270
Asp Tyr Phe Asp Pro His Gly Met His Ile Ala Gly Ile Leu Ala Gly					
	275		280		285
Asn Asp Thr Glu Gln Asp Ile Lys Asn Phe Asn Gly Ile Asp Gly Ile					
	290		295		300
Ala Pro Asn Ala Gln Ile Phe Ser Tyr Lys Met Tyr Ser Asp Ala Gly					
	305		310		315
Ser Gly Phe Ala Gly Asp Glu Thr Met Phe His Ala Ile Glu Asp Ser					
	325		330		335
Ile Lys His Asn Val Asp Val Val Ser Val Ser Ser Gly Phe Thr Gly					
	340		345		350
Thr Gly Leu Val Gly Glu Lys Tyr Trp Gln Ala Ile Arg Ala Leu Arg					
	355		360		365
Lys Ala Gly Ile Pro Met Val Val Ala Thr Gly Asn Tyr Ala Thr Ser					
	370		375		380
Ala Ser Ser Ser Ser Trp Asp Leu Val Ala Asn Asn His Leu Lys Met					



385                                      390                                      395                                      400  
 Thr Asp Thr Gly Asn Val Thr Arg Thr Ala Ala His Glu Asp Ala Ile  
    405                                      410                                      415  
 Ala Val Ala Ser Ala Lys Asn Gln Thr Val Glu Phe Asp Lys Val Asn  
    420                                      425                                      430  
 Ile Gly Gly Glu Ser Phe Lys Tyr Arg Asn Ile Gly Ala Phe Phe Asp  
    435                                      440                                      445  
 Lys Ser Lys Ile Thr Thr Asn Glu Asp Gly Thr Lys Ala Pro Ser Lys  
    450                                      455                                      460  
 Leu Lys Phe Val Tyr Ile Gly Lys Gly Gln Asp Gln Asp Leu Ile Gly  
 465                                      470                                      475                                      480  
 Leu Asp Leu Arg Gly Lys Ile Ala Val Met Asp Arg Ile Tyr Thr Lys  
    485                                      490                                      495  
 Asp Leu Lys Asn Ala Phe Lys Lys Ala Met Asp Lys Gly Ala Arg Ala  
    500                                      505                                      510  
 Ile Met Val Val Asn Thr Val Asn Tyr Tyr Asn Arg Asp Asn Trp Thr  
    515                                      520                                      525  
 Glu Leu Pro Ala Met Gly Tyr Glu Ala Asp Glu Gly Thr Lys Ser Gln  
    530                                      535                                      540  
 Val Phe Ser Ile Ser Gly Asp Asp Gly Val Lys Leu Trp Asn Met Ile  
 545                                      550                                      555                                      560  
 Asn Pro Asp Lys Lys Thr Glu Val Lys Arg Asn Asn Lys Glu Asp Phe  
    565                                      570                                      575  
 Lys Asp Lys Leu Glu Gln Tyr Tyr Pro Ile Asp Met Glu Ser Phe Asn  
    580                                      585                                      590  
 Ser Asn Lys Pro Asn Val Gly Asp Glu Lys Glu Ile Asp Phe Lys Phe  
    595                                      600                                      605  
 Ala Pro Asp Thr Asp Lys Glu Leu Tyr Lys Glu Asp Ile Ile Val Pro  
    610                                      615                                      620  
 Ala Gly Ser Thr Ser Trp Gly Pro Arg Ile Asp Leu Leu Leu Lys Pro  
 625                                      630                                      635                                      640  
 Asp Val Ser Ala Pro Gly Lys Asn Ile Lys Ser Thr Leu Asn Val Ile

645

650

655

Asn Gly Lys Ser Thr Tyr Gly Tyr  
660

&lt;210&gt; 7

&lt;211&gt; 2319

&lt;212&gt; DNA

<213> *Streptococcus pneumoniae*

&lt;400&gt; 7

```

aagcttggcg agattgcaga atctaaattt aaaaatttag gaaatggaaa agagggtagt 60
ctaaaaaaag atacaactgg ggtagaacat catcatcaag aaaatgaaga gtctattaaa 120
gaaaaatcta gttttactat tgatagaaat atttcaacaa ttagagactt tgaaaaataa 180
gacttaaaga aactcattaa aaagaaattt agagaagttg atgattttac aagtgaaact 240
ggtaagagaa tggaggaata cgattataaa tacgatgata aaggaaatat aatagcctac 300
gatgatggga ctgatctaga atatgaaact gagaaacttg acgaaatcaa atcaaaaatt 360
tatggtgttc taagtccgtc taaagatgga cactttgaaa ttcttggaaa gataagtaat 420
gtttctaaaa atgccaaggt atattatggg aataactata aatctataga aatcaaagcg 480
accaagtatg atttccactc aaaaacgatg acatttgatc tatacgctaa tattaatgat 540
attgtggatg gattagcttt tgcaggagat atgagattat ttgttaaaga taatgatcag 600
aaaaaagctg aaattaaaat tagaatgcct gaaaaaatta aggaaactaa atcagaatat 660
ccctatgtat caagttatgg gaatgtcata gaattagggg aaggagatct ttcaaaaaac 720
aaaccagaca atttaactaa aatggaatct ggtaaaatct attctgattc agaaaaacaa 780
caatatctgt taaaggataa tatcattcta agaaaaggct atgcactaaa agtgactacc 840
tataatcctg gaaaaacgga tatgttagaa ggaaatggag tctatagcaa ggaagatata 900
gcaaaaatac aaaaggccaa tcctaatacta agagcccttt cagaaacaac aatttatgct 960
gatagtagaa atgttgaaga tggagaagt acccaatctg tattaatgtc ggctttggac 1020
ggctttaaca ttataaggta tcaagtgttt acatttaaaa tgaacgataa aggggaagct 1080
atcgataaag acggaaatct tgtgacagat tcttctaaac ttgtattatt tggtaaggat 1140
gataaagaat aactggaga ggataagttc aatgtagaag ctataaaaga agatggctcc 1200
atgttatttta ttgataccaa accagtaaac ctttcaatgg ataagaacta ctttaatcca 1260
tctaaatcta ataaaattta tgtacgaaat ccagaatttt atttaagagg taagatttct 1320
gataaggggtg gttttaactg ggaattgaga gttaatgaat cggttgtaga taattattta 1380
atctacggag atttacacat tgataacact agagatttta atattaagct gaatgttaa 1440
gacggtgaca tcatggactg gggaaatgaaa gactataaag caaacggatt tccagataag 1500
gtaacagata tggatggaaa tgtttatctt caaactggct atagcgattt gaatgctaaa 1560
gcagttggag tccactatca gtttttatat gataatgtta aaccgaagt aaacattgat 1620
cctaagggaa atactagtat cgaatatgct gatggaaaat ctgtagtctt taacatcaat 1680
gataaaagaa ataatggatt cgatggtgag attcaagaac aacatattta tataaatgga 1740
aaagaatata catcatttaa tgatattaaa caaataatag acaagacact aaacattaag 1800
attgttgtaa aagattttgc aagaaataca accgtaaaag aattcatttt aaataaagat 1860
acgggagagg taagtgaatt aaaacctcat agggtaactg tgaccattca aaatggaaaa 1920
gaaatgagtt caacgatagt gtcggaagaa gattttatct tacctgttta taagggtgaa 1980
ttagaaaaag gataccaatt tgatggttgg gaaatttctg gtttcgaagg taaaaaagac 2040
gctggctatg ttattaatct atcaaaagat acctttataa aacctgtatt caagaaaata 2100
gaggagaaaa aggaggaaga aaataaacct acttttgatg tatcgaaaaa gaaagataac 2160

```

ccacaagtaa accatagtc attaaatgaa agtcacagaa aagaggattt acaaagagaa 2220  
 gagcattcac aaaaatctga ttcaactaag gatgttacag ctacagttct tgataaaaaac 2280  
 aatatcagta gtaaatcaac tactaacaat cctaataag 2319

<210> 8

<211> 773

<212> PRT

<213> Streptococcus pneumoniae

<400> 8

Lys Leu Gly Glu Ile Ala Glu Ser Lys Phe Lys Asn Leu Gly Asn Gly  
 1 5 10 15

Lys Glu Gly Ser Leu Lys Lys Asp Thr Thr Gly Val Glu His His His  
 20 25 30

Gln Glu Asn Glu Glu Ser Ile Lys Glu Lys Ser Ser Phe Thr Ile Asp  
 35 40 45

Arg Asn Ile Ser Thr Ile Arg Asp Phe Glu Asn Lys Asp Leu Lys Lys  
 50 55 60

Leu Ile Lys Lys Lys Phe Arg Glu Val Asp Asp Phe Thr Ser Glu Thr  
 65 70 75 80

Gly Lys Arg Met Glu Glu Tyr Asp Tyr Lys Tyr Asp Asp Lys Gly Asn  
 85 90 95

Ile Ile Ala Tyr Asp Asp Gly Thr Asp Leu Glu Tyr Glu Thr Glu Lys  
 100 105 110

Leu Asp Glu Ile Lys Ser Lys Ile Tyr Gly Val Leu Ser Pro Ser Lys  
 115 120 125

Asp Gly His Phe Glu Ile Leu Gly Lys Ile Ser Asn Val Ser Lys Asn  
 130 135 140

Ala Lys Val Tyr Tyr Gly Asn Asn Tyr Lys Ser Ile Glu Ile Lys Ala  
 145 150 155 160

Thr Lys Tyr Asp Phe His Ser Lys Thr Met Thr Phe Asp Leu Tyr Ala  
 165 170 175

Asn Ile Asn Asp Ile Val Asp Gly Leu Ala Phe Ala Gly Asp Met Arg  
 180 185 190

Leu Phe Val Lys Asp Asn Asp Gln Lys Lys Ala Glu Ile Lys Ile Arg

195	200	205
Met Pro Glu Lys Ile Lys Glu Thr Lys Ser Glu Tyr Pro Tyr Val Ser		
210	215	220
Ser Tyr Gly Asn Val Ile Glu Leu Gly Glu Gly Asp Leu Ser Lys Asn		
225	230	235 240
Lys Pro Asp Asn Leu Thr Lys Met Glu Ser Gly Lys Ile Tyr Ser Asp		
	245	250 255
Ser Glu Lys Gln Gln Tyr Leu Leu Lys Asp Asn Ile Ile Leu Arg Lys		
	260	265 270
Gly Tyr Ala Leu Lys Val Thr Thr Tyr Asn Pro Gly Lys Thr Asp Met		
	275	280 285
Leu Glu Gly Asn Gly Val Tyr Ser Lys Glu Asp Ile Ala Lys Ile Gln		
	290	295 300
Lys Ala Asn Pro Asn Leu Arg Ala Leu Ser Glu Thr Thr Ile Tyr Ala		
	305	310 315 320
Asp Ser Arg Asn Val Glu Asp Gly Arg Ser Thr Gln Ser Val Leu Met		
	325	330 335
Ser Ala Leu Asp Gly Phe Asn Ile Ile Arg Tyr Gln Val Phe Thr Phe		
	340	345 350
Lys Met Asn Asp Lys Gly Glu Ala Ile Asp Lys Asp Gly Asn Leu Val		
	355	360 365
Thr Asp Ser Ser Lys Leu Val Leu Phe Gly Lys Asp Asp Lys Glu Tyr		
	370	375 380
Thr Gly Glu Asp Lys Phe Asn Val Glu Ala Ile Lys Glu Asp Gly Ser		
	385	390 395 400
Met Leu Phe Ile Asp Thr Lys Pro Val Asn Leu Ser Met Asp Lys Asn		
	405	410 415
Tyr Phe Asn Pro Ser Lys Ser Asn Lys Ile Tyr Val Arg Asn Pro Glu		
	420	425 430
Phe Tyr Leu Arg Gly Lys Ile Ser Asp Lys Gly Gly Phe Asn Trp Glu		
	435	440 445
Leu Arg Val Asn Glu Ser Val Val Asp Asn Tyr Leu Ile Tyr Gly Asp		

450	455	460
Leu His Ile Asp Asn Thr Arg Asp Phe Asn Ile Lys Leu Asn Val Lys		
465	470	475 480
Asp Gly Asp Ile Met Asp Trp Gly Met Lys Asp Tyr Lys Ala Asn Gly		
	485	490 495
Phe Pro Asp Lys Val Thr Asp Met Asp Gly Asn Val Tyr Leu Gln Thr		
	500	505 510
Gly Tyr Ser Asp Leu Asn Ala Lys Ala Val Gly Val His Tyr Gln Phe		
	515	520 525
Leu Tyr Asp Asn Val Lys Pro Glu Val Asn Ile Asp Pro Lys Gly Asn		
	530	535 540
Thr Ser Ile Glu Tyr Ala Asp Gly Lys Ser Val Val Phe Asn Ile Asn		
	545	550 555 560
Asp Lys Arg Asn Asn Gly Phe Asp Gly Glu Ile Gln Glu Gln His Ile		
	565	570 575
Tyr Ile Asn Gly Lys Glu Tyr Thr Ser Phe Asn Asp Ile Lys Gln Ile		
	580	585 590
Ile Asp Lys Thr Leu Asn Ile Lys Ile Val Val Lys Asp Phe Ala Arg		
	595	600 605
Asn Thr Thr Val Lys Glu Phe Ile Leu Asn Lys Asp Thr Gly Glu Val		
	610	615 620
Ser Glu Leu Lys Pro His Arg Val Thr Val Thr Ile Gln Asn Gly Lys		
	625	630 635 640
Glu Met Ser Ser Thr Ile Val Ser Glu Glu Asp Phe Ile Leu Pro Val		
	645	650 655
Tyr Lys Gly Glu Leu Glu Lys Gly Tyr Gln Phe Asp Gly Trp Glu Ile		
	660	665 670
Ser Gly Phe Glu Gly Lys Lys Asp Ala Gly Tyr Val Ile Asn Leu Ser		
	675	680 685
Lys Asp Thr Phe Ile Lys Pro Val Phe Lys Lys Ile Glu Glu Lys Lys		
	690	695 700
Glu Glu Glu Asn Lys Pro Thr Phe Asp Val Ser Lys Lys Lys Asp Asn		

[illegible]

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 December 2000 (21.12.2000)

PCT


(10) International Publication Number  
**WO 00/76540 A3**

- (51) International Patent Classification<sup>7</sup>: A61K 39/09, C07K 16/12, C12N 5/10, A61K 39/40, A61P 31/04 // C07K 14/315
- (72) Inventors: ADAMOU, John, E.; 20822 Shamrock Glen Circle, Germantown, MD 20874 (US). CHOI, Gil, H.; 11429 Potomac Oaks Drive, Rockville, MD 20850 (US).
- (21) International Application Number: PCT/US00/15925
- (74) Agents: GRANT, Alan, J. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).
- (22) International Filing Date: 9 June 2000 (09.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/138,453 10 June 1999 (10.06.1999) US
- (81) Designated States (national): AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

[Continued on next page]

(54) Title: STREPTOCOCCUS PNEUMONIAE PROTEINS AND VACCINES

(57) Abstract: The present invention relates to novel immunogenic polypeptides, and fragments thereof, and vaccines for the prevention and treatment of pneumococcal infection, especially by *Streptococcus pneumoniae*. The invention also relates to antibodies against the disclosed polypeptides, as well as vaccines containing said polypeptides and methods of disease prevention.



WO 00/76540 A3



patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:  
8 February 2001

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 00/15925

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K39/09 C07K16/12 C12N5/10 A61K39/40 A61P31/04  
//C07K14/315

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 18930 A (HUMAN GENOME SCIENCES INC ;CHOI GIL H (US); HROMOCKYJ ALEX (US); J) 7 May 1998 (1998-05-07) page 4, line 6 -page 26 example 2 100% identity 38-664(seq.id 6):1-627 (seq.id 216 page 92-93) claims --- -/--	1-4, 9-13, 15-22

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

8 December 2000

Date of mailing of the international search report

14/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Covone, M

# INTERNATIONAL SEARCH REPORT

Intern. Patent Application No

PCT/US 00/15925

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NAYAK A R ET AL: "A LIVE RECOMBINANT AVIRULENT ORAL SALMONELLA VACCINE EXPRESSING PNEUMOCOCCAL SURFACE PROTEIN A INDUCES PROTECTIVE RESPONSES AGAINST STREPTOCOCCUS PNEUMONIAE" INFECTION AND IMMUNITY, AMERICAN SOCIETY FOR MICROBIOLOGY. WASHINGTON, US, vol. 66, August 1998 (1998-08), pages 3744-3751, XP000918253  ISSN: 0019-9567  the whole document</p>	20-22
P,X	<p>WO 00 06738 A (HANIFFY SEAN BOSCO ; LE PAGE RICHARD WILLIAM FALLA (GB); WELLS JER) 10 February 2000 (2000-02-10)  100% identity 1-664(seq.id.6):1-664 (seq.id.3 page 41-42)  100% identity 1-773(seq.id.8):1313-2086 (seq.id.3 page 41-42)  claims  example 2</p>	1-22

**FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

**1. Claims: 1-4,13 (completely) 9-12,15-22 (partially)**

A vaccine comprising a polypeptide, including immunogenic fragments thereof, having the amino acid sequence at least 65% identical to the amino acid sequence of Seq.ID 6. An antiserum or isolated antibody that bind said polypeptide, and an engineered cell producing said antibody. Methods to treat and/or prevent Streptococcus infection by administering said polypeptide or antibody.

**2. Claims: 5-8, 14 (completely) 9-12,15-22 (partially)**

A vaccine comprising a polypeptide, including immunogenic fragments thereof, having the amino acid sequence at least 65% identical to the amino acid sequence of Seq.ID 8. An antiserum or isolated antibody that bind said polypeptide, and an engineered cell producing said antibody. Methods to treat and/or prevent Streptococcus infection by administering said polypeptidy or antibody.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/15925

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9818930 A	07-05-1998	AU 5194598 A	22-05-1998
		AU 6909098 A	22-05-1998
		EP 0942983 A	22-09-1999
		EP 0941335 A	15-09-1999
		WO 9818931 A	07-05-1998
<hr/>			
WO 0006738 A	10-02-2000	NONE	
<hr/>			